

**Unclassified**  
**Defense Health Agency**  
**Deputy Assistant Director – Medical**  
**Affairs**



**DoD and VA Vision Center of Excellence DHA Practice**  
**Recommendation:**  
**Evaluation and Management of Dizziness**  
**Associated with Traumatic Brain Injury**  
**(TBI) for Eye Care Providers**

*Edition: 1*  
*October 2023*



**Defense Health Agency Falls Church, Virginia  
Evaluation and Management of Dizziness  
Associated with Traumatic Brain Injury (TBI)  
For Eye Care Providers  
Version 1.0  
2023**

**Release Authority:** Dr. Paul Cordts, Deputy Assistant Director – Medical Affairs, Defense Health Agency

**Document is UNCLASSIFIED**

**Editors:** see Authors and Affiliations

**Support From:** The Defense Health Agency Vision Center of Excellence, Hearing Center of Excellence, Traumatic Brain Injury Center of Excellence, Department of Defense and Department of Veterans Affairs

**Retrieve From:** <https://info.health.mil/sites/hro/CC/PracticeRecommendations/Forms/AllItems.aspx>

## **DHA Practice Recommendation: Overview and Disclaimer**

DHA Practice Recommendations (PRs) are developed by experts using the best information available at the time of publication. In some instances, some recommendations are expert opinion provided to users in the absence of definitive, well-designed, and executed randomized control trials. DHA's PRs provide an authoritative source of carefully synthesized clinical information. They are intended to assist clinical care teams with real-time decision making based on best available evidence.

While the DHA sponsors this PR, its endorsement of the findings and recommendations are limited to validation of the expert opinion and compiled evidence of the sponsoring subject matter expert (SME) body. This PR should be used to augment the practitioner's best clinical judgment. It may not account for local or structural conditions (i.e., resourcing, staffing, equipment, or Health Protection Conditions) affecting clinical decision making in the field by the practitioner.

DHA PRs are separate and distinct from jointly developed Department of Veterans Affairs (VA)/DoD Clinical Practice Guidelines that are the product of rigorous systematic literature review and synthesis. In contrast, DHA PRs provide the MHS practitioner with a synopsis of relevant clinical evidence tailored to the military medicine setting and TRICARE beneficiary population.

DHA PRs provide standardized evidence-informed guidelines that MHS practitioners should refer to when addressing patients with specific clinical conditions. Clinical practitioners must be mindful of the emergence of supervening clinical evidence published in the academic press not yet incorporated into the guideline.

This guideline is not intended to define a standard of care and should not be construed as such, nor should it be interpreted as prescribing an exclusive course of management for said condition or disease process. Variations in practice will inevitably and appropriately occur when clinicians consider the needs of individual patients, available resources, and limitations unique to an institution or type of practice. Every healthcare professional using this guideline is responsible for evaluating the appropriateness of applying it in the setting of any particular clinical situation.

This guideline does not represent TRICARE policy. Further, inclusion of recommendations for specific testing and/or therapeutic interventions within this guide does not guarantee coverage in Private Sector Care. Additional information on current TRICARE benefits may be found at [www.tricare.mil](http://www.tricare.mil) or by contacting the regional TRICARE Managed Care Support Contractor.

## Table of Contents

**Algorithm ..... 2**

**Purpose ..... 3**

**Diagnosis ..... 3**

**Expanded Patient Medical History ..... 3**

**Medication History ..... 4**

**Ocular History and Subjective Symptoms..... 4**

**Supplemental Ocular Testing..... 5**

**Management Considerations..... 6**

**Medical Terminology Used to Describe Patient Symptoms of Dizziness ..... 6**

**ICD-10 Coding Guidance for Dizziness following TBI..... 6**

**Additional Resources ..... 7**

**References..... 7**

**Statement of Authorship ..... 8**


**Authors..... 8**

**Potential Conflicts of Interest..... 9**

Vision Center of Excellence 2023 1.0 PR	
05 JULY 23	Date of expiration – N/A
Previous Document Number – N/A	Supersede date – N/A

# Algorithm

The “Algorithm for the Evaluation and Management of Dizziness Associated with TBI” will assist eye care providers in evaluation, management, and interdisciplinary care referral of patients presenting with dizziness following a concussion or brain injury of any severity.



## **Purpose**

Dizziness is a commonly reported symptom following head injury, including all severity levels of traumatic brain injury (TBI). Dizziness is a broad term, often encompassing more specific symptoms of vertigo, disequilibrium, presyncope, and vague lightheadedness. Vertigo is often described as a sensation or illusion of movement (such as spinning, rotating, tilting, or rocking), and typically denotes peripheral or central vestibular dysfunction.<sup>1</sup>

Estimates of 30-81% of individuals after TBI of all severities, and as high as 98% of those seen acutely following blast-related concussion, may experience dizziness and balance dysfunction.<sup>2,3,4</sup> Such patients can present to an eye care provider because visual dysfunctions can co-present with vestibular abnormalities post-TBI. The frequency of oculomotor dysfunction in patients with vestibular symptoms after concussion is higher compared with concussed patients without vestibular problems, thus requiring that such patients be assessed more vigorously.<sup>5</sup> It is important to recognize that treatable causes for dizziness and vertigo such as BPPV (Benign Paroxysmal Positional Vertigo) and migraine are very common and should be considered. More serious cases, which require prompt evaluation and care, can also be present in this population. Clinical presentations of dizziness following head trauma are often further complicated by polytrauma, medications, and other common co-morbid conditions, making diagnosis difficult. Referrals to members of a multidisciplinary TBI team or individual specialty providers are often required.

Multidisciplinary referral partners can include but are not limited to neurology, neuro-ophthalmology, physical medicine and rehabilitation, otolaryngology, audiology, physical therapy, primary care, internal medicine, cardiology, mental health specialties, ophthalmology, and optometry.

## **Diagnosis**

This PR will assist eye care providers in evaluating patients presenting with dizziness or vertigo and confirmed history of concussion or TBI of any severity. In some cases of severe TBI, this PR may be applied at a later stage when the patient sufficiently recovers. Use of this PR should follow a comprehensive dilated-eye examination and general medical evaluation to exclude systemic conditions such as hypotension or presyncope.<sup>6</sup> It provides guidance on obtaining an expanded patient medical history followed by supplemental testing recommendations and specialty referrals for further evaluation and management of dizziness. This PR should not replace sound clinical judgment or standard practice when caring for a patient.

## **Expanded Patient Medical History**

A comprehensive assessment of injury history can assist in evaluating and managing the patient's health. Review all relevant available multidisciplinary prior examinations and testing to determine if the patient's dizziness symptoms have already been evaluated to avoid duplicate referrals, including:

- Prior eye examinations
- Neurology evaluation
- Audiology evaluation
- Otolaryngology evaluation
- Vestibular evaluation

Refer to Tables 1 and 2 for guidance on dizziness-related injury history evaluation. When questioning patients, always use patient-friendly terminology and be mindful that a brain injury may slow the processing of the question and the response.

<b>Table 1. Detailed Dizziness-related Injury History by an Eye Care Provider</b>			
<b>Review TBI Events<sup>7</sup></b>	<b>Associated Symptoms</b>	<b>Characteristics of Symptoms</b>	<b>Relevant Past Medical History</b>
<ul style="list-style-type: none"> <li>• Date of injury</li> <li>• Location of injury</li> <li>• Loss of consciousness</li> <li>• Type of injury (blunt, blast, penetrating, motor vehicle accident, stroke, combined mechanism)</li> </ul>	<ul style="list-style-type: none"> <li>• Aural pressure or pain</li> <li>• Hearing loss</li> <li>• Tinnitus</li> <li>• Diplopia</li> <li>• Oscillopsia</li> <li>• Headache</li> <li>• Confusion/disorientation</li> <li>• Other neurological symptoms</li> </ul>	<ul style="list-style-type: none"> <li>• Duration</li> <li>• Frequency</li> <li>• Onset (continuous vs. episodic)</li> <li>• Precipitating factors (positional or postural effects)</li> <li>• Relieving factors</li> </ul>	<ul style="list-style-type: none"> <li>• Cardiovascular disease</li> <li>• Ear disease or surgery</li> <li>• Headaches/migraines (pre- and post-TBI)</li> <li>• Orthostatic hypotension</li> <li>• Prior vertigo</li> <li>• Psychological health disorders</li> <li>• Sleep disorders</li> <li>• Stressors/anxiety</li> <li>• Substance abuse disorders (e.g., drugs or alcohol)</li> </ul>

## Medication History

Many prescribed medications can contribute to dizziness and need to be evaluated.<sup>8,9</sup> (The list below is not exhaustive.)

- Abortive agents for migraine or migraine-like headaches
- Analgesics
- Antidepressants
- Anti-epileptic medications
- Anti-fatigue stimulant medications
- Anti-hypertensive medications
- Anxiolytic medications
- Non-steroidal, anti-inflammatory medications
- Prophylactic headache medications
- Psychotropic medications
- Sleep disorder medications (sedative/hypnotic)

## Ocular History and Subjective Symptoms

Numerous visual/ocular deficits can contribute to a patient's perception of dizziness after TBI. Examples of such contributory factors include: heightened sensitivity to blur from uncorrected refractive error; refractive correction adaptation, especially with astigmatic lenses, bifocals, progressives or wrap-style lenses; binocular, accommodative or oculomotor dysfunctions; motion perceived from flashes or floaters, especially involving more central vision; loss of peripheral vision altering visual-spatial awareness; photophobia/glare sensitivity; difficulty with vision at night contributing to blur or worsening of binocular status due to reduction in cues to peripheral fusion. Given the variety of visual/ocular deficits that may contribute to symptoms of dizziness, the following history questions should be used holistically and not individually in guiding the course of the eye exam.

<b>Table 2. Dizziness-related TBI Ocular History Questions<sup>7</sup></b>
<b>Dizziness-related Ocular History and Subjective Symptom Questions</b>
Do you have balance problems or dizziness?
Do you bump into objects or walls while walking?
Do you have restricted or missing part(s) of your visual field?
Does turning or tilting your head improve your vision or help your dizziness?
Is your vision blurry at distance or near since your injury?
Does covering one eye improve your overall vision?
Have you experienced any double vision since your injury?

Do you get headaches or brow aches during or after reading? Does this occur more frequently now than before your injury?
How often do you get headaches? If yes, do they occur with an episode of dizziness?
Is your vision blurry after reading?
Have you noticed a change in your ability to read since your injury?
What and how many materials do you read daily?
How long can you read continuously before you need to stop? Is this less than before your injury?
Do you experience sensitivity to light?
Do you see flashing lights or floating spots?
Is your vision at night worse since your injury?
Have you ever been diagnosed with benign paroxysmal positional vertigo (BPPV)?
Is there anything that you currently use that improves your vision since your injury?

## Supplemental Ocular Testing

Supplemental ocular testing is indicated if the initial comprehensive eye examination, correction of refractive error, and other updated testing do not resolve complaints and dizziness symptoms. This testing should be performed with the patient's habitual spectacle prescription first and repeated as needed after refraction if different from the habitual (see Table 3).

<b>Table 3: Supplemental Testing Guidance for Refractive and Oculomotor Dysfunction Post-Initial Comprehensive Eye Examination<sup>7,8</sup></b>	
<b>Considerations While Testing for Oculomotor Dysfunction</b>	
<b>Signs/Symptoms</b>	Diplopia Dizziness Headache Lack of clarity of thought/slowed thinking Lightheadedness Nausea Nystagmus (clinical sign) Oscillopsia Pain/discomfort Visual blur
<b>Tests for Oculomotor Dysfunction</b>	
<b>Extraocular Muscles (EOMs)</b>	Ductions/Versions Presence of Nystagmus (Central/Peripheral)
<b>Ocular Alignment in Free Space</b>	Cover Test at distance and near considerations: <ul style="list-style-type: none"> <li>– Patient positioned correctly (head/neck straight, no abnormal head posture)</li> <li>– Careful evaluation of vertical misalignment (ask about subjective movement of target)</li> <li>– Comitancy Testing (at every new visit and as needed thereafter, especially if patient is symptomatic and/or reports symptoms in a specific gaze other than primary)</li> </ul>
<b>Near Point of Convergence (NPC)*</b>	Repeat 3-5 times looking for regression (i.e., NPC recedes with repeated testing)
<b>Vergence Ranges in Free Space</b>	Distance and near Horizontal and vertical
<b>Accommodative Testing*</b>	Accuracy of target fixation Amplitude Facility
<b>Saccades and Pursuits*</b>	Inaccurate, hypometric saccades, and/or saccadic pursuits <ul style="list-style-type: none"> <li>– Horizontal</li> <li>– Vertical</li> </ul>



\*Detailed instructions on how to properly perform testing for near point of convergence, accommodative testing, and saccades/pursuits can be found in the DHA VCE Clinical Recommendation for the Eye Care Provider: Screening for Oculomotor Dysfunctions Following Traumatic Brain Injury (TBI). [Click here](#) for access to download **Clinical Screening Tool**.

## Management Considerations

Manage oculomotor dysfunction with refractive correction, near addition, prism, or vision rehabilitation. In conjunction with ocular management, proceed with appropriate specialty referrals based upon presentation, findings, and specialty availability at the local facility. If vertigo and signs of vestibular dysfunction are present, refer to neurology, otolaryngology, audiology, or vestibular physical therapy for further testing. If oscillopsia, acquired nystagmus, cranial nerve palsy, or unexplained strabismus are present, refer to ophthalmology, neurology, or neuro-ophthalmology. If headaches are prominent, refer to neurology or primary care for further evaluation for migraine. If symptoms suggest lightheadedness or near syncope, primary care follow-up is indicated (for more guidance see [Eye and Vision Care Following Blast Exposure and/or Possible Traumatic Brain Injury](#)). Consider limited duty status options until reduction in symptoms or resolution of condition.

## Medical Terminology Used to Describe Patient Symptoms of Dizziness<sup>9,10</sup>

- **Disequilibrium:** The patient describes a sensation that they are not quite where they think they should be or that their orientation to the world is “off.”
- **Imbalance:** The patient reports difficulty and unsteadiness when standing or walking.
- **Lightheadedness:** The patient reports a vague feeling in the head as if becoming weightless or feeling disconnected from the environment.
- **Near syncope:** The patient reports a feeling of almost fainting.
- **Oscillopsia:** The patient describes an illusion that the world is jiggling.
- **Vertigo:** The patient reports a sensation of motion when no motion is present or reports an altered sensation of motion when motion occurs. It is usually described as spinning (typically rotary) but can also be translational, tilting, swaying, rocking, or a linear motion.

## ICD-10 Coding Guidance for Dizziness following TBI

According to the DOD Clinical Recommendation Assessment and Management of Dizziness and Visual Disturbances Following Concussion/Mild Traumatic Brain Injury: Guidance for the Primary Care Manager, any patient who has sustained any severity level of a TBI and has persistent vision symptoms beyond 15 days should be referred to an eye care provider for a comprehensive vision and sensorimotor examination. The following are dizziness-related ICD-10 diagnostic codes following TBI that can be used.

## Eye Care Provider Dizziness-Related Diagnostic Codes

- **H81.9:** Unspecified disorder of vestibular function
- **R42:** Dizziness and giddiness, which includes disequilibrium

## Primary Care manager (PCM) or Subspecialty Dizziness-Related Diagnostic Codes

- **H53.19:** Other subjective visual disturbance
- **H81.39:** Other peripheral vertigo
- **H81.4:** Vertigo of central origin (used for suspected diagnosis)
- **H81.49:** Vertigo of central origin, unspecified ear (add location code as 5<sup>th</sup> character)
- **169.998:** Cervical vertigo (used for confirmed diagnosis)

This recommendation is not a substitute for existing guidance or clinical judgment. As with all clinical

decisions, field and operational circumstances may require deviation from these recommendations.

## **Additional Resources**

Assessment and Management of Oculomotor Dysfunctions Associated with Traumatic Brain injury. Department of Defense Vision Center of Excellence website. Updated December 13, 2016. [health.mil/Military-Health-Topics/Centers-of-Excellence/VCE/Provider-Resources](https://health.mil/Military-Health-Topics/Centers-of-Excellence/VCE/Provider-Resources)

Clinical Recommendation for the Eye Care Provider: Screening for Oculomotor Dysfunctions Following Traumatic Brain Injury (TBI). Defense Health Agency Vision Center of Excellence. 2021. [health.mil/Military-Health-Topics/Centers-of-Excellence/VCE/Provider-Resources](https://health.mil/Military-Health-Topics/Centers-of-Excellence/VCE/Provider-Resources)

Eye and Vision Care Following Blast Exposure and/or Possible Traumatic Brain Injury. Department of Defense Vision Center of Excellence website. Updated November 24, 2015. [health.mil/Military-Health-Topics/Centers-of-Excellence/VCE/Provider-Resources](https://health.mil/Military-Health-Topics/Centers-of-Excellence/VCE/Provider-Resources)

Rehabilitation of patients with Visual Field Loss Associated with Traumatic or Acquired Brain Injury. Department of Defense Vision Center of Excellence website. Updated April 27, 2016. [health.mil/Military-Health-Topics/Centers-of-Excellence/VCE/Provider-Resources](https://health.mil/Military-Health-Topics/Centers-of-Excellence/VCE/Provider-Resources)

## **References**

1. Cheridan, N. Dizziness. Cleveland Clinic Center for Continuing Education. August 2010. Last reviewed 2017. Accessed December 16, 2020. [clevelandclinicmeded.com/medicalpubs/diseasemanagement/neurology/dizziness/](https://clevelandclinicmeded.com/medicalpubs/diseasemanagement/neurology/dizziness/)
2. Maskell F, Chiarelli P, Isles R. Dizziness after traumatic brain injury: overview and measurement in the clinical setting. *Brain Inj.* 2006;20(3):293-305.
3. Hoffer ME, Balaban C, Gottshall K et al. Blast exposure: vestibular consequences and associated characteristics. *Otol Neurotol.* 2010;31(2):232-236.
4. Kisilevski V, Podoshin L, Ben-David J, et al. Results of otovestibular tests in mild head injuries. *Int Tinnitus J.* 2001;7(2):118-121.
5. Chinn RN, Marusic S, Wang A et al. Vergence and Accommodation Deficits in Children and Adolescents with Vestibular Disorders. *Optom Vis Sci.* DOI: 10.1097/OPX.0000000000001963.
6. Muncie HL, Sirmans SM, James E. Dizziness: Approach to Evaluation and Management. *Am Fam Physician.* 2017;95(3):154-162.
7. Goodrich GL, Martinsen GL, Flyg HM, et al. Development of a mild traumatic brain injury-specific vision screening protocol: a Delphi study. *J Rehabil Res Dev.* 2013;50(6):757-768.
8. Mucha A, Collins MW, Elbin RJ, et al. A Brief Vestibular/Ocular Motor Screening (VOMS) assessment to evaluate concussions: preliminary findings. *Am J Sports Med.* 2014;42(10):2479-2486.
9. Clinical Recommendation: Assessment and management of dizziness associated with mild TBI. Department of Defense and Veterans Brain Injury Center. 2012. Updated March 2020. [TBI Center of Excellence Provider Resources](#)
10. Coding Guidance for Diagnosing Vestibular Disorders in the MHS. Department of Defense Hearing Center of Excellence. June 2020. [hearing.health.mil/For-Providers/Diagnostic-and-Coding-Guidance](https://hearing.health.mil/For-Providers/Diagnostic-and-Coding-Guidance)

## Statement of Authorship

All listed authors, as well as peer reviewers and contributors, conceptualized the document, reviewed and revised the document critically for important intellectual content, approved the final document submitted, and agreed to be accountable for all work aspects. The VCE Branch Chief, DHA (R&E) ([VCE](#)) will ensure questions related to accuracy or integrity of any part of the work are appropriately investigated and resolved.

<b>Authors</b>	
<b>Potential Conflicts of Interest</b>	
The authors declare no conflicts of interest.	
<b>Defense Health Agency - Vision Center of Excellence (VCE) Silver Spring, MD/Walter Reed National Military Medical Center, Bethesda, MD</b>	
David Eliason, MD Deputy Chief	Natalya Merezhinskaya, PhD Health Research Scientist
Mark Reynolds, COL, MC, MPH, USA Chief, Army Public Health Center	Andrew Morgenstern, OD (CTR) Optometry Subject Matter Expert
Felix Barker, OD, MS Associate Director for Research	Michael Pattison, CAPT (r), OD, MS, USN Optometry Subject Matter Expert
Kevin Bradley, MS (CTR) Clinical Studies Analyst	
<b>Peer-Reviewers and Contributors Defense Health Agency – Hearing Center of Excellence (HCE)</b>	
Amy Boudin-George, AuD, CCC-A Acting Branch Lead Clinical Care, Rehabilitation, and Restoration Branch Defense Health Agency SAMHS, Joint Base San Antonio San Antonio, TX	Karen H. Lambert, PT, DPT, NCS Vestibular Program Manager Contractor, zCore Business Solutions Clinical Care, Rehabilitation, and Restoration Section Defense Health Agency San Antonio, TX
Carlos Esquivel, MD, FACS Chief Medical Officer, HCE Research and Engineering, Defense Health Agency Wilford Hall Ambulatory Surgery Center Joint Base San Antonio San Antonio, TX	Seyi Gbade-Alabi, MAJ, MD, MC, FAAPMR, USA Assistant Professor & Director, Undergraduate Medical Education Department of PM&R USUHS School of Medicine Bethesda, MD
<b>Defense Health Agency – Traumatic Brain Injury Center of Excellence (TBICoE)</b>	
Faith Akin, PhD Professor Department of Audiology & Speech Pathology East Tennessee University Johnson City, TN	Scott William Pyne, CAPT (r), MD, MC, USN Assistant Professor, Department of Family Medicine, USUHS Family and Sports Medicine Physician United States Naval Academy Naval Health Clinic Annapolis, MD
Gary McKinney, MS, DHSc, CBIS Chief, Office of Clinical Practice and Clinical Recommendations Research and Engineering, Defense Health Agency Silver Spring, MD	Terryl Aitken, OD Fort Belvoir Community Hospital Fort Belvoir, VA

<b>Authors</b>	
<b>Potential Conflicts of Interest</b>	
The authors declare no conflicts of interest.	
Katharine Stout, DPT, NCS, MBA Assistant Branch Chief, TBICoE Research and Engineering, Defense Health Agency Silver Spring, MD	
<b>Department of Defense</b>	
Beverly Scott, COL (r), MC, USA Medical Advisory Board Seattle, WA	Marcy Pape, PT, DPT, CLT, CBIS, CCVT Physical Therapist National Intrepid Center of Excellence (NICoE) Walter Reed National Military Medical Center Bethesda, MD
Geeta Girdher, OD Optometrist, National Intrepid Center of Excellence (NICoE) Director, Residency & Student Externship Programs Walter Reed National Military Medical Center Bethesda, MD	Nathan T. Tagg, COL (r), MC, USA Associate Professor of Ophthalmology and Neurology Department Duke University School of Medicine Durham, NC
<b>Department of Veterans Affairs</b>	
Sally H. Dang, OD, MPH Chief, Optometry Service VA Long Beach Healthcare System Long Beach, CA	Randy Kardon, MD, PhD Director, Center for the Prevention and Treatment of Visual Loss University of Iowa Iowa City, IA
Chrystyna Rakoczy, OD (Working Group Chair) Clinic Director Brain Injury Vision Assessment and Rehabilitation/Polytrauma-TBI Optometry-PREP James A. Haley Veterans' Hospital Tampa, FL	Suzanne Wickum, OD (Working Group Chair) Staff Optometrist Tibor Rubin VA Medical Center Long Beach, CA
<b>Intrepid Spirit Center</b>	
Robin Winslow, OD Neuro-Optometric Vision Rehabilitation Intrepid Spirit Center Fort Belvoir Community Hospital Fort Belvoir, VA	

Approved By  
**CORDTS.PAUL.RO**  
 GER.1044303619

Digitally signed by  
 CORDTS.PAUL.ROGER.1044303619  
 Date: 2023.07.05 10:45:40 -04'00'

**5 July 2023**

**Signature**

**Date of Signature**

Paul R. Cordts, MD  
 Senior Executive Service  
 Deputy Assistant Director, Medical Affairs  
 Defense Health Agency